

I 49.89:82-47-1/map C 11

VALDOSTA, GEORGIA–FLORIDA

30 X 60 MINUTE SERIES (TOPOGRAPHIC)

MAP C11

Valdosta

FLORIDA ECOLOGICAL ATLAS

SOILS AND LANDFORMS



MMS 85-0011
FWS/OBS-82/47.1



ORIGINAL

Scale 1:100 000

TOPICAL MAPS IN THIS ATLAS:

A. BIOLOGICAL RESOURCES
B. SOCIOECONOMIC FEATURES
C. SOILS AND LANDFORMS
D. OIL, GAS AND MINERALS
E. HYDROLOGY AND CLIMATOLOGY

1984

Produced by U.S. Fish and Wildlife Service and Minerals Management Service

Projection and 10,000 meter grid, zone 17, Universal



Produced by U.S. Fish and Wildlife Service and Minerals Management Service

Projection and 10,000 meter grid, zone 17, Universal Transverse Mercator
25,000-foot grid ticks based on Florida coordinate system, west zone

Overlay material compiled in 1983 from latest available data obtained from Federal, State and Local government agencies and other sources.

MAJOR SURFACE LANDFORMS

Western Highlands	Fountain Slopes
Escambia Valley	Gravel Ridge
Gulf Coastal Lowlands	Northern Highlands
Gulf Barrier Chain	Coastal Swamps
Wicahema Lowlands	Bradenville Ridge
New Hope Ridge	Western Valley
Greenhead Slopes	

SOIL ASSOCIATIONS

1. Lakeland-Kanlo Lakewood Association
2. Lakeland, Red Gulfport Sand Somerset, Lakewood, Red Gulfport Association
3. Red Bay-Wakeley Association
4. Tifton-Carnegie Fuquay Association
5. Norfolk-Ruston-Savannah Association
6. Orangeburg-Ruston-Wakeley Association
7. Klej Lean Association
8. Hockaden-Kalmia-Ungela Association
9. Coastal Dune Land and Beach-Tidal Marsh Association
10. Pioneer-Ruffage-Alluvial Land Fresh Water Swamp-Alluvial Land Tidal Marsh
11. Lakeland-Pasta-Kureb-St. Lucie-Remin Association
12. Lakeland-Troup Association
13. Troup-Orangeburg Association
14. Dothan-Tarpon-Ona Association


NORTHERN HIGHLANDS
THOMAS CO
GEORGIA
FLORIDA
JEFFERSON CO
BROOKS CO
MADISON CO
HAMILTON CO
Lake Miccosukee
Tallahassee
Thomasville
Quitman
Valdosta
Monticello
Madison
Live Oak
Jasper
White Springs
Lake City
Perry
Chiefland
Old Town
Cross City
Steinhatchee
Branford
Jennings
Fargo
Homerville
Nahunta
Waycross
Folkston
Fernandina Beach
Jacksonville
Gainesville
Ocala
Inverness
Brooksville
Tarpon Springs
Dade City
Zephyrhills
Tampa
Plant City
Lakeland
Winter Haven
Bartow
Sebring
Avon Park
Lake Placid
Arcadia
Wauchula
Fort Meade
Bradenton
Sarasota
Venice
Punta Gorda
Fort Myers
Naples
Everglades City
Marco Island
Key West
Miami
Homestead
Florida City
Key Largo
Tavernier
Islamorada
Marathon
Big Pine Key
Lower Matecumbe Key
Upper Matecumbe Key
Long Key
Duck Key
Grassy Key
Summerland Key
Cudjoe Key
Sugarloaf Key
Boca Chica Key
Stock Island
Tallahassee Int'l Airport
Turkey Point
Homestead AFB
Eglin AFB
Duke Field
Camp Blanding
Cecil Field
Jacksonville NAS
Mayport NAS
Kennedy Space Center
Patrick AFB
Orlando Int'l Airport
MacDill AFB
Tampa Int'l Airport
Albany
Moultrie
Tifton
Douglas
Fitzgerald
Cordele
Vidalia
Swainsboro
Statesboro
Brunswick
Waycross
Valdosta
Thomasville
Bainbridge
Dothan
Ozark
Enterprise
Andalusia
Opp
Crestview
Fort Walton Beach
Destin
Panama City
Apalachicola
Carrabelle
Port St. Joe
Wewahitchka
Blountstown
Marianna
Chipley
Bonifay
DeFuniak Springs
Milton
Pensacola
Gulf Breeze
Perdido Key
Lake Seminole
Lake Talquin
Lake Miccosukee
Lake Iamonia
Lake Jackson
Lake Lafayette
Lake Miccosukee
Lake Rousseau
Lake Tsala Apopka
Lake Panasoffkee
Lake Okeechobee
Lake Kissimmee
Lake Istokpoga
Lake Placid
Lake Jackson
Lake Okeechobee
Lake Tohopekaliga
Lake Hatchineha

SCALE 1:100 000

1 CENTIMETER ON THE MAP REPRESENTS 1 KILOMETER ON THE GROUND
CONTOUR INTERVAL 5 METERS

KILOMETERS 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20
MILES 0 1 2 3 4 5 6 7 8 9 10 11 12

1000 0 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000 11000 12000 13000 14000 15000 16000 17000 18000 19000 20000 METERS
5000 0 5000 10000 15000 20000 25000 30000 35000 40000 45000 50000 55000 60000 65000 FEET

VALDOSTA, GEORGIA–FLORIDA

N3030–W8300/30 X 60

1981

THE FLORIDA ECOLOGICAL ATLAS WAS PRODUCED BY THE U.S. FISH AND WILDLIFE SERVICE, DIVISION OF BIOLOGICAL SERVICES, NATIONAL COASTAL ECOSYSTEMS TEAM, SLIDELL, LOUISIANA. THE ATLAS WAS DESIGNED TO ASSIST FEDERAL AGENCY PLANNERS, DECISION MAKERS, AND THE GENERAL PUBLIC, TO DETERMINE AND EVALUATE CHANGE AND TO RESOLVE CONFLICTS AMONG LAND USES, FISH AND WILDLIFE RESOURCES, AND HABITAT PROTECTION. THE GEOGRAPHIC AREA OF THE STUDY AREA INCLUDES THE COASTAL COUNTIES: ESCAMBIA, SANTA ROSA, OKALOOSA, WALTON, BAY, GULF, CALHOUN, GULF, FRANKLIN, WAKULLA, JEFFERSON, TAYLOR, DIXIE, LEVY, CITRUS, HERNANDO, PASCO, PINELLAS, HILLSBOROUGH, MANATEE, SARASOTA, CHARLOTTE, LEE, COLLIER, AND DADE COUNTIES IN FLORIDA. THE ATLAS DESIGN AND PRODUCTS OF THIS STUDY ARE NOT INTENDED TO BE USED TO ESTABLISH JURISDICTION OF ANY FEDERAL, STATE, OR LOCAL GOVERNMENT TO ESTABLISH REGULATORY TECHNICAL SCOPE OF THE REGULATORY PROGRAMS OF GOVERNMENT AGENCIES.

THIS MAP HAS BEEN COMPILED FROM A VARIETY OF SOURCE MATERIALS INCLUDING MAPS, AERIAL PHOTOS, AND REPORTS. SINCE MANY SOURCES CONTAINED GENERALIZED LOCATIONS OF RESOURCES AND OTHER MAPPED FEATURES, THE USER SHOULD NOT RELY ON THESE MAPS FOR PRECISE MEASUREMENTS OF DETAILS OF VALUE.

PRIMARY SOURCES OF MAPPED INFORMATION:

LANDFORMS: Photographs Map of Florida, State Bureau of Geology, 1982

SOILS: General Soils Atlas of Florida, Soil Conservation Service, 1981
General Soils Atlas of Florida, U.S. Soil Conservation Service, 1982

ACTIVE DUNES: Black and White Aerial Photographs, 1:80,000 Scale, Mark Hurd Productions, Inc., 1985
Topographic Maps, 1:24,000 Scale, U.S. Geological Survey, various dates

GEOLOGIC FAULTS: Map of Escambia and Santa Rosa Counties Showing Contours on top of the Pollock Limestone Member of the Apalachicola Formation, Florida Bureau of Geology, 1985

FOR COMPLETE REFERENCES SEE MAP NARRATIVE

10. Plummer-Sutlage-Alhonial Land-Fresh Water Swamp Duncan-Pamlico Association: Nearly level, poorly drained soils with a thick surface layer overlying a loamy subsoil and very poorly drained soils that are sandy throughout.

19. Fresno-Leon-Troup Association: Nearly level to sloping, well drained soils with sandy loams over loamy subsoil.

23. Ardilla-Lenfield-Orlinon Association: Nearly level to gently sloping, somewhat poorly drained and moderately well drained sandy soils with loamy subsoil.

44. Dothan-Orangeburg-Red Bay Association: Nearly level to sloping well drained sandy or loamy soils with loamy subsoils.

E=__ ft/yr — BEACH EROSION AREA (Period of Record: 1957-1981; Univ. of West Florida, 1981)

A=__ ft/yr — BEACH ACCRETION AREA (Period of Record: 1957-1981; Univ. of West Florida, 1981)

GULF BARRIER ISLAND

ACTIVE DUNE AREA

U D — NORMAL FAULT (Showing relative vertical movement; U. up, and D. down)

U D — SUB-SURFACE NORMAL FAULT (Showing relative vertical movement; U. up, and D. down)

STUDY AREA BOUNDARY

Offshore

Inland (County)

Inland (State)



1. Chipley-Blanton-Swamp Association
2. Albemarle, var. Chiefland-Blanton Association
3. Cardin-Apopka Association
4. Lowndesville-Chiefland Association
5. Fernandina-Centenary-Otela Association
6. Orangeburg-Ruston-Red Bay Association
7. Spirit-Lochloosa-Tavares Association
8. Albemarle-Osier Association
9. Albany-Lucy-Centenary Association
10. Dothan-Thomasville-Kenansville Association
11. Leon-Wacasha-Surrency Association
12. Lynne-Fortune-Pamlapex Association
13. Cardin-Paola-Tavares Association
14. Albany-Centenary-Lucy Association
15. Arredonda-Sandrock Association
16. Brewer-Bexa Association
17. Fernandina-Bulfield Association
18. Wadley-Bonnegar Association
19. Fresno-Leon-Troup Association
20. Terra Ceia-Pluvial Association
21. Arredonda-Spen-Kendrick Association
22. Cardin-Lake Association
23. Wasnet Association
24. Albany-Bonnegar-Orrington Association
25. Eustis-Wekiwa-Sunnyside Association
26. Paisley-Flordana-Wabasso Association
27. Blanton-Arapaha-Terra Ceia Association
28. Hontoosa-Weekenetney-Lissochee Association
29. Felda-Immokalee Association
30. Salt Water Marsh and Swamp Association
31. Fresh Water Swamp and Marsh Association
32. Bulfield-Fernandina Association
33. Paisley-Bulfield Association
34. Wadley-Penella Association

REPRODUCED FROM BEST AVAILABLE COPY